Student Introduction

Daniel Jumper University of Illinois at Urbana-Champaign Spinfest 2014

Introduction

From:

Dallas, Texas



• Undergrad:

- Abilene Christian University ('05-'09)
- Worked on PHENIX summer '06, '07


• Grad School:

- University of Illinois at Urbana-Champaign ('09present) under Matthias Grosse Perdekamp
- Work:
 - Local UIUC RPC r&d
 - RPC assembly(St. 1) and operation (Run 13)
 - Run 13 $W \rightarrow \mu A_L$ analysis

Slideshow!







RPC 3 "Efficiency" Hodoscope

RPC 1
Before/After
Installation



Slideshow!



RPC 1 Before/After Installation



My Analysis: Run 13 $W \rightarrow \mu A_L$

Goal:

• Better constrain separate q and \bar{q} contributions to longitudinal proton spin

Channel:


- $q\bar{q} \rightarrow W^{\pm} \rightarrow \mu \nu_{\mu}$ (at forward rapidity)
- Weak interaction: parity violation
 - Constrains possible quark helicities giving a simplified $A_L \sim \Delta q$, $\Delta \bar{q}$ relation


$$A_L^{W^+} = \frac{-\Delta u(x_1)\bar{d}(x_2) + \Delta \bar{d}(x_1)u(x_2)}{u(x_1)\bar{d}(x_2) + \bar{d}(x_1)u(x_2)} \quad A_L^{W^-} = \frac{-\Delta d(x_1)\bar{u}(x_2) + \Delta \bar{u}(x_1)d(x_2)}{d(x_1)\bar{u}(x_2) + \bar{u}(x_1)d(x_2)}$$

My Analysis: Run 13 $W \rightarrow \mu A_L$

- Challenge
 - Background dominance!
- Technique
 - Likelihood ratio (W_{ness})
 - Use signal/background differences in many variables to select a subset of data with more relative signal
 - Maximum Likelihood Signal/Background fit
 - Characterize remaining sig/bkg ratio to correct asymmetries

My Analysis: Run 13 $W \rightarrow \mu A_L$





Bonus: Claim to fame

PHENIX Themeo

First cake featured in the elog

http://logbook2.phenix.bnl.gov:7815/Run+13+Log/3323

